

WHAT IS CLAIMED IS:

- 1 1. A method of detecting cancer cells in a biological sample from a
2 mammal, the method comprising the steps of:
 - 3 (i) providing the biological sample from the mammal; and
 - 4 (ii) detecting an overexpression of a Pellino 1 polypeptide comprising at least
5 70% amino acid identity to SEQ ID NO:2 or a Pellino 2 polypeptide comprising at least 70%
6 amino acid identity to SEQ ID NO:4 in the biological sample, thereby detecting the presence
7 of cancer cells in the biological sample.
- 1 2. The method of claim 1, wherein the Pellino 1 polypeptide has an
2 amino acid sequence of SEQ ID NO:2 or the Pellino 2 polypeptide has an amino acid
3 sequence of SEQ ID NO:4.
- 1 3. The method of claim 1, wherein the Pellino 1 or Pellino 2 polypeptide
2 is detected using an antibody that selectively binds to the polypeptide.
- 1 4. The method of claim 1, wherein the detecting step comprises detecting
2 an mRNA that encodes the Pellino 1 or Pellino 2 polypeptide.
- 1 5. The method of claim 1, wherein the cancer cells are from an epithelial
2 cancer.
- 1 6. The method of claim 5, wherein the epithelial cancer is a lung, colon,
2 or ovarian cancer.
- 1 7. The method of claim 1, wherein the mammal is a human.
- 1 8. A method of detecting cancer cells in a biological sample from a
2 mammal, the method comprising the steps of:
 - 3 (i) providing the biological sample from the mammal; and
 - 4 (ii) detecting an increase in copy number of a gene encoding a Pellino 1
5 polypeptide comprising at least 70% amino acid identity to SEQ ID NO:2 or a Pellino 2
6 polypeptide comprising at least 70% amino acid identity to SEQ ID NO:4 in the biological
7 sample, thereby detecting the presence of cancer cells in the biological sample.
- 1 9. The method of claim 8, wherein the detecting step further comprises:

(a) contacting the gene with a probe that selectively hybridizes to the gene under conditions in which the probe selectively hybridizes to the gene to form a stable hybridization complex; and

(b) detecting the hybridization complex.

10. The method of claim 8, wherein the Pellino 1 polypeptide has an amino acid sequence of SEQ ID NO:2 or the Pellino 2 polypeptide has an amino acid sequence of SEQ ID NO:4.

11. The method of claim 8, wherein the cancer cells are from an epithelial cancer.

12. The method of claim 11, wherein the epithelial cancer is a lung, colon, or ovarian cancer.

13. The method of claim 8, wherein the mammal is a human.

14. A method of monitoring the efficacy of a therapeutic treatment of cancer, the method comprising the steps of:

(i) providing a biological sample from a mammal undergoing the therapeutic treatment; and

(ii) detecting a level of a Pellino 1 polypeptide comprising at least 70% amino acid identity to SEQ ID NO:2 or a Pellino 2 polypeptide comprising at least 70% amino acid identity to SEQ ID NO:4, or detecting copy number of a gene encoding the Pellino 1 or Pellino 2 polypeptide in the biological sample compared to a level or copy number in a biological sample from the mammal prior to, or earlier in, the therapeutic treatment; thereby monitoring the efficacy of the therapy.

15. The method of claim 14, wherein the Pellino 1 polypeptide has an amino acid sequence of SEQ ID NO:2 or the Pellino 2 polypeptide has an amino acid sequence of SEQ ID NO:4.

16. The method of claim 14, wherein the cancer is an epithelial cancer.

17. The method of claim 15, wherein the epithelial cancer is a lung, colon, or ovarian cancer.

- 1 18. The method of claim 14, wherein the mammal is a human.
- 1 19. A method of identifying a compound that inhibits the activity of a
2 Pellino 2 polypeptide, the method comprising the steps of:
3 (i) contacting the compound with a Pellino 2 polypeptide that comprises at
4 least 90% identity to an amino acid sequence of SEQ ID NO:4; and
5 (ii) detecting a decrease in the activity of the Pellino 2.
- 1 20. The method of claim 19, wherein the Pellino 1 polypeptide has an
2 amino acid sequence of SEQ ID NO:2 or the Pellino 2 polypeptide has an amino acid
3 sequence of SEQ ID NO:4.
- 1 21. The method of claim 19, wherein the Pellino 2 polypeptide is
2 amplified in the cell compared to normal.
- 1 22. A method of inhibiting proliferation of a cancer cell that
2 overexpresses a Pellino 1 polypeptide comprising at least 70% amino acid identity to SEQ
3 ID NO:2 or a Pellino 2 polypeptide comprising at least 70% amino acid identity to SEQ ID
4 NO:4, the method comprising the step of contacting the cancer cell with a therapeutically
5 effective amount of an inhibitor of the Pellino 1 or 2 polypeptide.
- 1 23. The method of claim 22, wherein the cancer cell is from an epithelial
2 cancer.
- 1 24. The method of claim 23, wherein the epithelial cancer is a lung, colon,
2 or ovarian cancer cell.
- 1 25. The method of claim 22, wherein the Pellino 1 polypeptide has an
2 amino acid sequence of SEQ ID NO:2 or the Pellino 2 polypeptide has an amino acid
3 sequence of SEQ ID NO:4.
- 1 26. The method of claim 22, wherein the inhibitor is identified using the
2 method of claim 19.
- 1 27. The method of claim 22, wherein the inhibitor is an antibody.

1 28. The method of claim 22, wherein the inhibitor is an antisense
2 polynucleotide.

1 29. A isolated nucleic acid encoding a Pellino 2 polypeptide, wherein the
2 Pellino 2 polypeptide comprises at least 95% amino acid sequence identity to SEQ ID NO:4.

1 30. The nucleic acid of claim 29, wherein the nucleic acid encodes a
2 Pellino 2 polypeptide comprising an amino acid sequence of SEQ ID NO:4.

1 31. The nucleic acid of claim 29, wherein the nucleic acid comprises a
2 nucleotide sequence of SEQ ID NO:3.

1 32. An expression vector comprising the nucleic acid of claim 29.

1 33. A host cell comprising the expression vector of claim 32.

1 34. An isolated Pellino 2 polypeptide comprising at least 95% amino
2 acid identity to SEQ ID NO:4.

1 35. The polypeptide of claim 34, wherein the polypeptide comprises an
2 amino acid sequence of SEQ ID NO:4.

1 36. The polypeptide of claim 34, wherein the polypeptide specifically
2 binds to antibodies generated against a polypeptide comprising an amino acid sequence of
3 SEQ ID NO:4.

1 37. An antibody that specifically binds to the polypeptide of claim 34.